



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Allovahlkampfia minuta nov. sp., (Acrasidae, Heterolobosea, Excavata) a new soil amoeba at the boundary of the acrasid cellular slime moulds.

Citation for published version:

Alvaro DE OBESO FERNANDEZ DEL VALLE, A & Maciver, S 2017, 'Allovahlkampfia minuta nov. sp., (Acrasidae, Heterolobosea, Excavata) a new soil amoeba at the boundary of the acrasid cellular slime moulds.', *Acta Protozoologica*, vol. 56, no. 3. <https://doi.org/10.4467/16890027AP.17.016.7497>

Digital Object Identifier (DOI):

[10.4467/16890027AP.17.016.7497](https://doi.org/10.4467/16890027AP.17.016.7497)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Acta Protozoologica

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Allovahlkampfia minuta nov. sp., (Acrasidae, Heterolobosea, Excavata) a
new soil amoeba at the boundary of the acrasid cellular slime moulds.

Alvaro DE OBESO FERNADEZ DEL VALLE, Sutherland K MACIVER*

Biomedical Sciences, Edinburgh Medical School, University of Edinburgh, Hugh Robson
Building, George Square. Edinburgh. EH8 9XD. Scotland. UK.

*Corresponding author. Phone 44(0)131 650 3714, E-mail smaciver@ed.ac.uk

Page heading: *Allovahlkampfia minuta*, a new heterolobosean

17

18 **ABSTRACT**

19 We report the isolation of a new species of *Allovahlkampfia*, a small cyst-forming
20 heterolobosean soil amoeba. Phylogenetic analysis of the 18S rDNA and the internal
21 transcribed spacers indicates that *Allovahlkampfia* is more closely related to the acrasids than
22 to other heterolobosean groups and indicates that the new strain (GF1) groups with
23 *Allovahlkampfia tibetiensis* and *A. nederlandiensis* despite being significantly smaller than
24 these and any other described *Allovahlkampfia* species. GF1 forms aggregated cyst masses
25 similar to the early stages of *Acrasis* sorocarp development, in agreement with the view that it
26 shares ancestry with the acrasids. Time-lapse video microscopy reveals that trophozoites are
27 attracted to individuals that have already begun to encyst or that have formed cysts. Although
28 some members of the genus are known to be pathogenic the strain GF1 does not grow above
29 28°C nor at elevated osmotic conditions, indicating that it is unlikely to be a pathogen.

30

31 **INTRODUCTION**

32 The class heterolobosea was first created on morphological grounds to unite the schizopyrenid
33 amoebae/amoebflagellates with the acrasid slime moulds (Page & Blanton, 1985), and
34 subsequent molecular genetic data supported this union (Roger *et al.* 1996; Keeling and
35 Doolittle, 1996). The locomotory habit of these amoebae is to tend to produce an eruptive
36 pseudopod, often alternatively from one side and then the other, of the advancing front. The
37 heterolobosea are not directly related to other amoebae such as those within the Amoebozoa,
38 despite being similar in appearance and habit. The heterolobosean acrasid slime moulds are
39 very similar to the amoebozoan slime moulds too in life cycle, but these remarkable similarities
40 in appearance and function are most probably due to parallel evolution.

It is not usually possible to classify the trophozoites of this group morphologically since they share very similar habits and appearances but many heterolobosean amoebae are also capable of transforming into flagellates, these are so called amoeboflagellates. In addition, some heterolobosean groups are comprised of flagellates only (e.g. *Percolomonas* and *Stephanopogon*) having apparently lost the ability to adopt the ancestral amoeboid habit. Heterolobosean amoebae which do not form flagellates were traditionally placed in the genus *Vahlkampfia* (Page 1976) but it has since been shown that this group is not monophyletic and other genera such as *Paravahlkampfia* and *Neovahlkampfia* were formed (Brown and De Jonckheere 1999), to join other non-flagellate genera raised on morphological grounds. Further analysis of 18S RNA genes has allowed many other organisms to be included with the Heterolobosea and it is increasingly clear that it is an order with huge diversity (Pánek and Čepička 2016; Pánek *et al.* 2017). In 2009 it was necessary to raise a further genus, named *Allovahlkampfia* to place a new heterolobosean that did not fit elsewhere (Walochnik and Mulec 2009). A number of other isolates have now been added and as the genus *Allovahlkampfia* is presently configured, does not contain flagellates or amoeboflagellates but all members so far characterised, produce cysts without opercula. Members of *Allovahlkampfia* vary significantly in SSU rRNA sequence, with inserts present in some (Geisen *et al.* 2015). They are variable in size but the strain described in the present study which we name *Allovahlkampfia minuta*, is the smallest.

One heterolobosean amoeboflagellate, *Naegleria fowleri* is dangerously pathogenic for humans (Carter 1970; Siddiqui *et al.* 2016). Additionally, *Allovahlkampfia spelaea* is reported to infect the human eye (Huseein *et al.* 2016; Tolba *et al.* 2016) and to host bacteria known to be human pathogens (Mohamed *et al.* 2016) making this genus particularly worthy of study.

METHODS AND MATERIALS

Soil samples were collected from the Glenfinnan viaduct at the north end of Loch Shiel, Scotland, UK. Small quantities were settled on non-nutrient agar overlain with Neff's saline overnight. Strips of this were then inverted onto non-nutrient agar plates spread with *E. coli* (BL21) and monitored over a week for amoebal out growth. Small blocks were cut from the leading outgrowth onto fresh agar plates spread with *E. coli* until clones of strain GF1 were established. Large quantities of cloned GF1 amoebae were cultured on 2% agar YME plates containing 0.01% yeast extract and 0.025% malt extract overlain with Neff's saline at room temperature with *E. coli* as food organisms. For long term storage, the amoebae and cysts used in this study were taken up in Neff's saline with 10% DMSO and placed overnight in a freezer at -20°C, then transferred to -80°C freezer. Amoebae were also stored long term in clay pellets as previously described (Lorenzo-Morales and Maciver 2006).

To test tolerance to osmotic stress, amoebae were seeded on *E. coli* spread non-nutrient agar plates supplemented with 0.5M, 1M and 1.5M mannitol as previously described (Khan *et al.* 2001). Temperature tolerance was also tested by seeding amoebae on *E. coli* spread non-nutrient agar plates incubated at various temperature and monitored for growth over a ten day period.

We investigated the ability of *A. minuta* to develop a flagellate stage with a variety of stimuli included the addition of distilled water, rapid changes in pH, CO₂ and O₂ levels in addition to physical agitation (Perkins and Jahn 1970).

DNA was isolated from GF1 as previously described (Lorenzo-Morales *et al.* 2005). Amoebae were lysed in buffer and treated with proteinase K at 60°C for 2 hours followed by phenol-chloroform extraction. DNA was then concentrated by precipitation with isopropanol. The resulting DNA was quantified using a "nanodrop" spectrophotometer (ThermoFisher). An

aliquot of 100ng of genomic DNA was used per PCR reaction, using GoTaq Green Master Mix polymerase (Promega).

Sequences were obtained from GenBank and other sources and compiled together with the new sequences from this study using “Seaview”, version 4 (Gouy *et al.* 2010). “BioEdit” (Hall 1999) was used to trim sequences and to determine levels of homology between sequences. Sequences were aligned using seaview4 which was also used to implement the PhyML algorithm to produce Maximum likelihood phylogenetic trees (Guindon and Gascuel 2003) using the GTR model. The non-parametric analysis was performed with 100 bootstrap pseudo-replicates, using representative sequences from other heterolobozoans (*Naegleria*, *Tetramitus* and *Vahlkampfia*) as the outgroup. Binomial names of the strains described previously (Geisen *et al.* 2015) were obtained from the CCAP website (<https://www.ccap.ac.uk>).

RESULTS

The locomotive form is noticeably smaller than that of previously described members of this genus (table 1). It is often wider than it is long although this is usually temporary. Pseudopods are produced eruptively from the anterior in a manner typical of the heterolobosean amoebae (Fig. 1). The trophozoites can also adopt a flabellate-like morphology. In culture GF1 trophozoites and cysts were observed to collect at the water air boundary and migrated on the meniscus. Strain GF1 was found to produce aggregates of cysts on agar plates (Fig. 2). Time-lapse video microscopy (see supplementary data and Fig. 3) reveals that the aggregated nature of the cyst distribution results from trophozoites actively migrating toward preformed cysts and not as a result of cyst aggregating by a passive mechanism after formation. GF1 was found to grow at 4°C and slowly at 28°C but at 30°C it encysts. The optimal temperature for growth was 24°C. GF1 could not grow on plate supplemented with 0.5M mannitol or above.

Phylogenetic analysis tentatively suggests that genus *Allovahlkampfia* is comprised of eight or nine groups, probably representing as many natural species. More data from more genes and isolates are required to clarify this of course. No flagellated cells have been found in culture despite the application of a range of previously reported flagellate inducing stimuli. Strain GF1 is closely related to two previously reported species (*A. nederlandiensis* and *A. tibetiensis*) (Fig. 4a) and these are very similar in appearance also, especially regarding the tendency to adopt a temporary flabellate locomotory habit. However, these other members of this group are significantly larger (Geisen *et al.* 2015) (Table 1).

DISCUSSION

The most notable feature of *A. minuta* is its small size but similarly sized heterolobosean amoebae have been described. For example, *Neovahlkampfia nana* is 12 µm in length (Tymł *et al.* 2017). Compared to other heteroloboseans including the genus *Allovahlkampfia*, the locomotion of *A. minuta* seems chaotic and inefficient with frequent direction changes and the production of seemingly immediately redundant pseudopods. *A. minuta* was observed to adopt a temporal flabellate morphology as is the case for other heteroloboseans for example *Heteramoeba clara* (Droop 1962) and *Vahlkampfia signyensis* (Garstecki *et al.* 2005). The habit of the cysts and trophozoites of *A. minuta* to collect at the meniscus has been reported to be the case with other free-living amoebae (Preston 2003).

The genus *Allovahlkampfia* is genetically diverse but morphologically uniform and represented so far only by freshwater and soil species. It is not possible to distinguish *Allovahlkampfia* from several other heterolobosean genera based on appearance alone (Geisen *et al.* 2015), even at the level of electron microscopy (González-Robles *et al.* 2012). The cysts of all presently named species of the genus *Allovahlkampfia*, including *A. minuta* described here, lack opercula (Anderson *et al.* 2011; Walochnik and Mulec 2009; Geisen *et al.* 2015). This is a feature shared

139 by *Acrasis rosea* (Hohl and Hamamoto 1969). It is the small size of *A. minuta* that
140 distinguishes it from others in the genus. None of the previously characterised members of the
141 genus is close in size to the trophozoites or the cysts.

142 Since reports of morphological similarities between vahlkampfid and acrasids were published
143 (Page 1978; Page and Blanton 1985), several authors have reported a close genetic affinity also
144 (Keeling and Doolittle 1996; Roger *et al.* 1996). Comparing ribosomal genes allowed
145 *Allovahlkampfia* and *Acrasis* to be placed in clade “Acrasidia” within the Tetramitida (De
146 Jonckheere *et al.* 2011; Harding *et al.* 2013; Geisen *et al.* 2015; Pánek *et al.* 2017). In this
147 connection, the Allovahlkampfid strain AND12 (Lara *et al.* 2007) is especially interesting as it
148 is currently the most closely known relative of the genus *Acrasis*. However, it was only
149 partially characterized and because it was observed in liquid culture, its ability to develop
150 fruiting bodies is untested (Lara *et al.* 2007). Unfortunately, this strain is no longer available
151 in culture (Ekelund, Pers. Comm), but amoebae at the boundary between *Allovahlkampfia* and
152 *Acrasis* may reveal details on the development of sporogenesis.

153 The aggregation of *A. minuta* trophozoites during encystment, produces images very similar to
154 those seen in the acrasids especially the morphologically simple *Acrasis helenhemmesae* (see
155 figures 14-16 in Brown *et al.* 2010). This tendency to aggregate may be shared with others of
156 the genus as *A. palustris* and *A. spelaea* cysts are also reported to occur in clumps, although no
157 further information about this was given (Anderson *et al.* 2011; Tolba *et al.* 2016). The
158 significance of this aggregation is not yet clear but we have found time lapse video evidence
159 (see supplementary data) for chemotaxis towards the encysting centre. There may be increased
160 survival of aggregated cysts, or the aggregation may, like is this case in the acrasids be a prelude
161 to the production of Allovahlkampfid fruiting bodies that may form under as-yet unknown
162 circumstances. In this regard, it is interesting to note that a simple linear fruiting body has been
163 observed in *Allovahlkampfia* strain BA (Brown *et al.* 2012), however this was observed on

only one occasion. Other hints that *Allovahlkampfia* can produce multicellular aggregates such as fruiting exist. The isolate, *Allovahlkampfia* strain OSA, has been described as “an orange stalk-like fungal growth habit” (Shutt 2006) and while this description is a little vague, it is compatible with it having a multi-cellular or at least aggregative stage. Confusingly however, neither BA nor OSA formed fruiting bodies according to the worker who isolated them (Schutt 2006). Strains BA and OSA were assigned to *A. spelaea* (Brown *et al.* 2012) but this suggestion has not been adopted by later workers (Geisen *et al.* 2015) and is not supported by our 18S rDNA analysis. The aggregation that we observed with encysting *A. minuta* amoeba is superficially similar to the aggregation during the encystment of the amoebozoan, *Vannella pentlandii* (Maciver *et al.* 2017). Also, the closely related *Vannella fimicola* produces an actual fruiting body (Olive, 1962). The ability to fruit has evolved apparently independently, in other amoebozoan groups, including *Luapeleamoeba hula* within family Acanthamoebidae (Shadwick *et al.* 2016), and surprisingly, including the genus *Acanthamoeba* itself (Tice *et al.* 2016). Amoebae other than the Amoebozoa are also known to produce fruiting bodies (Kang *et al.* 2017). There is little evidence of sexual recombination in the acrasids (Brown *et al.* 2012), but it is likely that the purpose of these fruiting bodies in the acrasids may be a dispersive strategy carrying the spores off the substrate to be carried by air currents, or passing birds and insects.

Many genera of free living amoeba contain species pathogenic to humans and while most of these including *Acanthamoeba*, *Balamuthia* and *Sappinia* are members of the amoebozoa some are heteroloboseans. *Acanthamoeba* is notorious for causing *Acanthamoeba* keratitis (Lorenzo-Morales *et al.* 2015), a sight threatening and extremely painful infection of the eye but there are reported cases of a similar keratitis associated with the presence of amoebae tentatively identified as being *Vahlkampfia* but in the presence of other organisms (Aitken *et al.* 1996; Alexandrakis *et al.* 1998; Niyyati *et al.* 2010; González-Robles *et al.* 2012). These

reports, while not concluding a directly causal link, do reveal that that heteroloboseans can survive on the human eye. Stronger evidence suggests that an amoeba SO/1P now known to belong in genus *Allovahlkampfia* (Fig. 2), cause keratitis in humans with herpes virus (Ozkoc *et al.* 2008) and that this same strain produced a similar infection in rat models (Huseein *et al.* 2016) (presumably in the absence of herpes virus since it was not mentioned). A second case where an amoeba closely related to *Allovahlkampfia spelaea* was reported to cause keratitis has also been reported (Tolba *et al.* 2016) and this strain also caused keratitis in a rabbit model. Whereas 93% of *Acanthamoeba* keratitis cases are associated with contact lens use (Radford *et al.* 1998), the few cases reported for heterolobosean amoebae have been dominated by trauma to the eye surface (Alexandrakis *et al.* 1998; Ozkoc *et al.* 2008; Tolba *et al.* 2016). It is firmly established that another heterolobosean, *Naegleria fowleri* causes the usually fatal brain infection primary amoebic meningoencephalitis in humans (Siddiqui *et al.* 2016). The finding that *A. minuta* does not grow above 28°C, and its failure to cope with elevated osmolarity, would seem to rule it out as a potential human eye pathogen where the temperature varies between 32 and 36°C (Purslow *et al.* 2005).

Allovahlkampfia minuta n. sp.

Description. The locomotive morphology of this species is typical for small heterolobosean amoebae. Amoeba length 13.4 µm (range 9.5 – 16.7) and width 8.1 µm (range 4.6 – 15.6). Length:breadth ratio is 1.65. A well-developed uroid is not often visible but trailing uroidal filaments are occasionally observed on glass and tissue culture plastic. The spherical nucleus is variable in diameter but it ranges between 1.5 - 2.5 µm. There is a central nucleolus. A contractile vacuole is usually visible in trophozoites in Neff's saline or distilled water. This is usually located at the rear of the amoeba, often within the bulbous uroid, if it is present. Growth

on *E. coli* is observed between 4 and 28°C, with an optimum at 24°C. The pelleted trophozoites have a distinct pink/orange colour.

Cysts. Spheroidal diameter 6.5 µm (range 4.7 – 8.9) with a single cell wall and lacking pores.

The mononuclear cysts are usually formed in aggregates.

Etymology. This isolate is significantly smaller than any other described *Allovahlkampfia* species, hence the name “*minuta*”.

Type locality. Bank of Loch Shiel (freshwater) near the Glennfinnan monument (latitude 56.8687; longitude -5.4378), Scotland, UK. The elevation is 4m.

Habitat. Alluvial soil.

Type material. A culture of *A. minuta* has been deposited with the Culture Collection of Algae and Protists, accession number CCAP 2502/7, and gene sequences deposited with GenBank, MF680037 and MF677901 for the 18S rRNA gene and the ITS region respectively.

ACKNOWLEDGEMENTS

We are grateful to Mr Samuel Casasola Zamora for collecting soil samples. We are pleased to acknowledge support from Mexico's National Council of Science and Technology (CONACYT) and a University of Edinburgh College of Medicine and Veterinary Medicine PhD Studentship. Support from Amoebics Ltd, Edinburgh is also acknowledged. We also gratefully acknowledge two anonymous reviewers for their useful and insightful comments on an earlier version of the manuscript.

REFERENCES

Aitken D., Hay J., Kinnear F.B., Kirkness C.M., Lee W.R., Seal D.V. (1996) Amebic keratitis in a wearer of disposable contact lenses due to a mixed *Vahlkampfia* and *Hartmanella* infection. *Ophthalmol.* **103**: 485-494

241 Anderson O.R., Wang W.E.N., Faucher S.P., Bi K., Shuman H.A. (2011) A new
 242 heterolobosean amoeba *Solumitrus palustris* n. g., n. sp. isolated from freshwater marsh
 243 soil. *J. Eukaryot. Microbiol.* **58**: 60–67
 244
 245 Alexandrakis G., Miller D., Huang A.J.W. (1998) Amebic keratitis due to *Vahlkampfia*
 246 infection following corneal trauma. *Arch. Ophthalmol.* **117**: 950-951
 247
 248 Brown S., De Jonckheere J.F. (1999) A reevaluation of the amoeba genus *Vahlkampfia* based
 249 on SSUrDNA sequences. *Eur. J. Protistol.* **35**: 49-54
 250
 251 Brown M.W., Silberman J.D., Spiegel F.W. (2010) A morphologically simple species of
 252 *Acrasis* (Heterolobosea, Excavata), *Acrasis helenhemmesae* n. sp. *J. Eukaryot. Microbiol.*
 253 **57**: 346–353
 254
 255 Brown M.W., Silberman J.D., Spiegel F.W. (2012) A contemporary evaluation of the acrasids
 256 (Acrasidae, Heterolobosea, Excavata). *Eur. J. Protistol.* **48**: 103–123
 257
 258 Carter R.F. (1970) Description of a *Naegleria* species isolated from two cases of primary
 259 amoebic meningoencephalitis and of the experimental pathological changes induced by it.
 260 *J. Pathol.* **100**: 217–244
 261
 262 De Jonckheere J.F., Murase J., Opperdoes F.R. (2011) A new thermophilic heterolobosean
 263 amoeba, *Fumarolamoeba ceborucoi*, gen. nov., sp. nov., isolated near a fumarole at a
 264 volcano in Mexico. *Acta Protozool.* **50**: 41–48
 265

266 Droop M. R. (1962) *Heteramoeba clara* n. gen., n. sp., a sexual biphasic amoeba. *Arch.*
267 *Microbiol.* **42**: 254–266
268

269 Garstecki T., Brown S., De Jonckheere J.F. (2005) Description of *Vahlkampfia signyensis* n.
270 sp. (Heterolobosea), based on morphological, ultrastructural and molecular characteristics.
271 *Eur. J. Protistol.* **41**: 119–127
272

273 Geisen S., Bonkowski M., Zhang J., De Jonckheere J.F. (2015) Heterogeneity in the genus
274 *Allovalkampfia* and the description of the new genus *Parafumarolamoeba*
275 (*Vahlkampfiidae*; Heterolobosea). *Eur. J. Protistol.* **51**: 335–349
276

277 González-Robles A., Salazar-Villatoro L., González-Lázaro M., Omaña-Molina M., Martínez-
278 Palomo A. (2012) *Vahlkampfia* sp: Structural observations of cultured trophozoites. *Exp.*
279 *Parasitol.* **130**: 86–90
280

281 Gouy M., Guindon S., Gascuel O. (2010) SeaView Version 4: A multiplatform graphical user
282 interface for sequence alignment and phylogenetic tree building. *Mol. Biol. Evol.* **27**: 221-
283 224
284

285 Guindon S., Gascuel O., (2003) A simple, fast, and accurate algorithm to estimate large
286 phylogenies by maximum likelihood. *Syst. Biol.* **52**: 696–704
287

288 Hall T.A., (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis
289 program for Windows 95/98/NT. *Nucl. Acids Symp. Ser.* **41**: 95–98
290

291 Harding T., Brown M.W., Plotnikov A., Selivanova E., Park J.S., Gunderson J.H., Baumgartner
292 M., Silberman J.D., Roger A.J., Simpson A.G. (2013) Amoeba stages in the deepest
293 branching heteroloboseans, including *Pharyngomonas*: evolutionary and systematic
294 implications. *Protist* **164**: 272–286
295
296 Hohl H.R., Hamamoto S.T. (1969) Ultrastructure of *Acrasis rosea*, a cellular slime mold,
297 during development. *J. Protozool.* **16**: 333-344
298
299 Huseein E.A.M., Anwer F.E.S., Abed G.H., Abdel-Gahfar K.S., Omar H.E.D.M. (2016)
300 Keratitis induced by *Allovahlkampfia spelaea* in experimental rat model: a trial for
301 treatment with ellagic acid. *Eur. J. Biol. Res.* **6**: 299-309
302
303 Kang S., Tice A.K., Spiegel F.W., Silberman J.D., Pánek T., Čepička I., Kostka M., Kosakyan
304 A., Alcântara D.M., Roger A.J., Shadwick L.L., Smirnov A., Kudryavstev A., Lahr D.J.G.,
305 Brown M.W. (2017) Between a pod and a hard test: the deep evolution of amoebae. *Mol.*
306 *Biol.Evol.* **34**: 2258–2270
307
308 Keeling P.J., Doolittle W.F. (1996). Alpha-tubulin from early diverging eukaryotic lineages
309 and the evolution of the tubulin family. *Mol. Biol. Evol.* **13**: 1297–1305
310
311 Khan N.A., Jarroll E.J., Paget, T.A. (2001) *Acanthamoeba* can be differentiated by the
312 polymerase chain reaction and simple plating assays. *Curr Microbiol.* **43**: 204–208
313

314 Lara E., Berney C., Ekelund F., Harms H., Chatzinotas A. (2007) Molecular comparison of
 315 cultivable protozoa from a pristine and a polycyclic aromatic hydrocarbon polluted site.
 316 *Soil Biol.Biochem.* **39**: 139–148
 317

318 Lorenzo-Morales J., Lindo J.F., Martinez E., Calder D., Figueruelo E., Valladares B., Ortega-
 319 Rivas A. (2005) Pathogenic *Acanthamoeba* strains from water sources in Jamaica, West
 320 Indies. *Annals Trop. Med. & Parasitol.* **99**: 751–758
 321

322 Lorenzo-Morales J., Maciver S.K. (2006) Long-term, low maintenance storage of
 323 *Acanthamoeba* strains. *Parasitology Research* **99**: 743-745
 324

325 Lorenzo-Morales J., Khan N.A., Walochnik, J. (2015) An update on *Acanthamoeba* keratitis:
 326 diagnosis, pathogenesis and treatment. *Parasite* **22**: 10-20
 327

328 Maciver S.K., De Obeso Fernandez Del Valle A., Koutsogiannis Z. (2017) *Vannella pentlandii*
 329 n. sp., (Amoebozoa, Discosea, Vannellida) a small, cyst-forming soil amoeba. *Exp.*
 330 *Parasitol. (in press)*
 331

332 Mohamed M.E., Huseein E.A., Farrag H.M., Mostafa F.A.A., Hassan A.T. (2016)
 333 *Allovahlkampfia spelaea* is a potential environmental host for pathogenic bacteria. *J.*
 334 *Bacteriol. Parasitol.* **7**: 255-262
 335

336 Niyyati M., Lorenzo-Morales J., Rezaie S., Rahimi F., Martín-Navarro C.M., Mohebalı M.,
 337 Rezaeian M. (2010) First report of a mixed infection due to *Acanthamoeba* genotype T3
 338 and *Vahlkampfia* in a cosmetic soft contact lens wearer in Iran. *Exp. Parasitol.* **126**: 89-90

339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363

Olive L. S. (1962) The genus *Protostelium*. *Amer. J. Botany* **49**: 297-303

Ozkoc S., Tuncay S., Delibas S. B., Akisu C., Ozbek Z., Durak I., Walochnik J. (2008) Identification of *Acanthamoeba* genotype T4 and *Paravahlkampfia* sp. from two clinical samples. *J. Med. Microbiol.* **57**: 392–396

Page F.C. (1976) An illustrated key to freshwater and soil amoebae. Freshwater Biological Association, Ambleside, England.

Page F.C. (1978) *Acrasis rosea* and the possible relationship between Acrasida and Schizopyrenida. *Archiv für Protistenkunde.* **120**: 169-181

Page F.C., Blanton R.L. (1985) The Heterolobosea (Sarcodina:Rhizopoda), a new class uniting the Schizopyrenida and the Acrasidae (Acrasida). *Protistologica* **21**: 121–132

Pánek T., Čepička I. (2016) Diversity of Heterolobosea. In “Genetic Diversity of Microorganisms”, pp. 3–26. Eds. Caliskan M., Rijeka M : InTech.

Pánek T., Simpson A.G.B., Brown M.W., Dyer B.D. (2017) Heterolobosea in “Handboook of the Protists” Archibald *et al.*, eds Springer Int. Publ.

Perkins D.L., Jahn T.L. (1970) Amoeboflagellate transformations and the Gibbs-Donnan ratio. *J. Protozool.* **17**: 168-172

364 Preston T.M. (2003) The water-air interface: a microhabitat for amoebae. *Eur. J. Protistol.* **39**:
 365 385–389
 366
 367 Purslow C., Wolffsohn J.S., Santodomingo-Rubido J. (2005) The effect of contact lens wear
 368 on dynamic ocular surface temperature. *Contact Lens & Ant. Eye* **28**: 29-36
 369
 370 Radford C.F., Lehmann O.J., Dart J.K. (1998) *Acanthamoeba* keratitis: multicentre survey in
 371 England 1992–1996. *Br J Ophthalmol.* **82**: 1387–1392
 372
 373 Roger A.J., Smith M.W., Doolittle R.F., Doolittle W.F. (1996) Evidence for the heterolobosea
 374 from phylogenetic analysis of genes encoding glyceraldehyde-3-phosphate
 375 dehydrogenase. *J. Euk. Microbiol.* **43**: 475-485
 376
 377 Shadwick L.L., Brown M.W., Tice A.K., Spiegel F.W. (2016) A new amoeba with Protosteloid
 378 Fruiting: *Luapeleamoeba hula* n. g. n. sp. (Acanthamoebidae, Centramoebida,
 379 Amoebozoa) *Acta Protozool.* **55**: 123–134
 380
 381 Shutt T.E. (2006) Origins and evolution of mitochondrial transcription and replication
 382 machinery. Ph.D. Thesis, Dalhousie University, Halifax, Nova Scotia.
 383
 384 Siddiqui R., Ali K.M., Cope J.R., Khan N.A. (2016) Biology and pathogenesis of *Naegleria*
 385 *fowleri*. *Acta Tropica* **164**: 375–394
 386
 387 Tice A.K., Shadwick L.L., Fiore-Donno A.M., Geisen S., Kang S., Schuler G.A., Spiegel F.W.,
 388 Wilkinson K.A., Bonkowski M., Dumack K., Lahr D.J.G., Voelcker E., Clauß S., Zhang

J. Brown M.W. (2016) Expansion of the molecular and morphological diversity of Acanthamoebidae (Centramoebida, Amoebozoa) and identification of a novel life cycle type within the group. *Biology Direct* **11**: 69 DOI 10.1186/s13062-016-0171-0


Tolba M.E.M., Huseein E.A.M., Farrag H.M.M., Mohamed H.E.D., Kobayashi S., Suzuki J., Sugano S. (2016) *Allovahlkampfia spelaea* causing keratitis in humans. *PLoS Negl Trop Dis.* **10**: e0004841

Tymł T., Skulinová K., Kavan J., Ditrich O., Kostka M., Dyková I. (2016) Heterolobosean amoebae from Arctic and Antarctic extremes: 18 novel strains of *Allovahlkampfia*, *Vahlkampfia* and *Naegleria*. *Eur. J. Protistol.* **56**: 119-133

Tymł T., Lares-Jiménez L.F., Kostka M., Dyková I. (2017) *Neovahlkampfia nana* n. sp. Reinforcing an underrepresented subclade of Tetramitida, Heterolobosea. *J. Euk. Microbiol.* **64**: 78–87

Walochnik J., Mulec, J. (2009) Free-living amoebae in carbonate precipitating microhabitats of karst caves and a new vahlkampfiid amoeba, *Allovahlkampfia spelaea* gen. nov., sp. nov. *Acta Protozool.* **48**: 25–33


414 **Figures and Legends**



415

416 Figure 1

417 **A)** Trophozoites of *Allovahlkampfia minuta* showing typical morphology with eruptive
418 pseudopods. Nucleus (N) has a central nucleolus and the contractile vacuole (CV) is typically
419 positioned at the rear of the amoeba. The scale bar is 20 µm. **B)** Cysts. The scale bar is 10 µm.
420 Food organisms, *E. coli* are visible in the background.



421

Figure 2

Trophozoites gather and encyst on non-nutrient agar plate surface as the *E. coli* food organisms become depleted and as the conditions become less moist.





Figure 3

Cooperative cyst formation. **A)** Trophozoites were placed on an agar plate with a monolayer of *E. coli*. Encystment took place when the plate became dry and the food source locally exhausted. Trophozoites aggregated around the first to encyst, forming a slightly raised mound of cysts. **B)** This image is the first frame of a video (see supplementary data) taken to follow cyst aggregate formation. **C)** This is the last image of the same video where black arrowheads point to cysts that have been produced during the time that the video was shot.



B). A PhyML phylogenetic tree (GTR model) of members of some heteroloboseans of the genus *Allovahlkampfia* and *Acrasis* based on the fragment coding the internal transcribed spacer 1, 5.8S ribosomal RNA, and internal transcribed spacer 2.

Table 1

Species/strain	Length μm Average &/or range	Breadth μm Average & range	Cyst diameter μm Average &/or range	Reference
<i>A. palustris</i>	31 (14 – 43)	6.3 (4 – 12)	8 – 10	Anderson <i>et al.</i> 2011
<i>A. spelaea</i>	20 – 40	n/a	16 – 25	Walochnik and Mulec, 2009
AND12	20 – 30	n/a	n/a	Lara <i>et al.</i> 2007
<i>A. tibetiensis</i>	25.2 (21.2–31.2)	6.6 (5.6–9.2)	6.6 (5.0–8.2)	Geisen <i>et al.</i> 2015
<i>A. nederlandiensis</i>	33.2 (21.8 – 50.8)	5.8 (3.6 – 8.6)	6.2 (5.0- 7.8)	Geisen <i>et al.</i> 2015
F11	35.8	11.4	11.5	Tyml <i>et al.</i> 2016
<i>A. minuta</i>	13.4 (9.5 – 16.7)	8.1 (4.6 – 15.6)	6.5 (4.7 – 8.9)	This study

Supplementary data

Video 1. Locomotion of *Allovahlkampfia minuta* trophozoites. Real time video micrography shows typical eruptive locomotion but also show the chaotic nature of the progression of this species.

Video 2. Cyst chemotaxis. Time lapse video with frames every 15 seconds, total time 3 hours. While some trophozoites continue to move feed and divide, others are drawn toward existing cysts and they too then encyst. Contractile vacuole activity gradually lessens as cyst development takes place.